

CONNECTION BETWEEN DIFFRACTION AND SMALL-X

N.N. NIKOLAEV

*Institut f. Kernphysik, Forschungszentrum Jülich
D-52425 Jülich, Germany*

and

*L.D. Landau Institute for Theoretical Physics
142432 Chernogolovka, Russia*

E-mail: N.Nikolaev@fz-juelich.de

A brief review of the modern QCD theory of diffractive DIS is given.

1 Why diffractive DIS is so fundamental

Let us dream of e-Uranium DIS at THERA at $Q^2 \sim 10 \text{ GeV}^2$ and $x \sim 10^{-5}$. Violent DIS is associated with complete destruction of the target, deposition of a mere dozen MeV energy breaks the uranium nucleus entirely, yet a rigorous prediction from unitarity is that *diffractive* DIS $eU \rightarrow e' XU$ with the target nucleus emerging intact in the ground state will make $\approx 50\%$ of total DIS ¹! By the Glauber-Gribov theory, the abundance of diffraction is closely related to nuclear shadowing (NS) in DIS. In 1974 Nikolaev and Zakharov reinterpreted NS in terms of the saturation of nuclear parton densities ². More recently, the NZ picture of saturation has been addressed to within QCD, see Iancu's talk at this Symposium ³. If correct, this QCD approach must inevitably lead to 50 % diffractive DIS. To summarize, diffractive DIS is a key to nuclear parton densities and QCD predictions for the initial state in ultrarelativistic nuclear collisions. Because at HERA the rate of diffractive DIS is mere 10%, saturation effects are all but marginal, see ⁴ and talks ⁵ at this Symposium .

2 Color dipole link between inclusive and diffractive DIS

SF's of DIS are related by optical theorem to the forward virtual Compton scattering (CS), which in the color dipole (CD) factorization ⁶ takes the form $A_{CS} = \Psi_f^* \otimes A_{q\bar{q}} \otimes \Psi_{in}$ where $\Psi_{f,in}$ is the wave function (WF) of the $q\bar{q}$ Fock state of the photon and the $q\bar{q}$ -proton scattering kernel $A_{q\bar{q}}$ is proportional to the color dipole cross section, which for small dipoles is related to the gluon SF of the target, $\sigma(x, \mathbf{r}) \approx \frac{\pi^2}{3} r^2 \alpha_S(\frac{A}{r^2}) G(x, \frac{A}{r^2})$. where $A \approx 10$ by properties of the Bessel functions ⁷. Taking for Ψ_f^* the WF of the vector meson (VM) or the $X = q\bar{q}$ plane waves gives the diffraction excitation of VM ⁴ or hadronic




Figure 1. The test of the $(Q^2 + m_V^2)$ scaling. The divergence of the solid and dashed curves indicates the sensitivity to the WF of the VM. The experimental data are from HERA ¹⁴.

continuum, $\gamma^* p \rightarrow X p'$. Generalization to excitation of the $q\bar{q}g$ or higher Fock states of the photon is straightforward ^{8,9}, with certain reservations CD results can be reinterpreted in terms of the parton structure of the pomeron with the solid result ^{8,9,10} that gluons and charged partons carry about an equal fraction of the momentum of the pomeron. The formalism set in ^{8,9} and especially in ^{10,11} is a basis of modern parameterizations of the diffractive structure function ¹². Unfortunately the use of the discredited Ingelmann-Schlein-Regge factorization and DGLAP evolution which is not warranted at large β ^{9,11}, in the still simplified form of this analysis makes conclusions ^{12,13} on the gluon content of the pomeron highly suspect.

3 The $Q^2 + m_V^2$ scaling

While DIS probes CD $\sigma(x, \mathbf{r})$ in a broad range of $\frac{1}{AQ^2} \lesssim r^2 \lesssim 1 \text{ fm}^2$, the diffractive VM production probes $\sigma(x, \mathbf{r})$ at a scanning radius ^{8,4} $r \sim r_S =$




Figure 2. The $(Q^2 + m_V^2)$ scaling of the effective intercept and diffraction slope ¹⁷

$6/\sqrt{Q^2 + m_V^2}$, and the gluon SF of the target at the hard scale $\bar{Q}^2 \approx (0.1-0.25)*(Q^2 + m_V^2)$ and $x = 0.5(Q^2 + m_V^2)/(Q^2 + W^2)$. After factoring out the charge-isospin factors, that entails the $(Q^2 + m_V^2)$ scaling of the VM production cross section⁴, see fig. 1. The same scaling holds also for the effective intercept $\alpha_{\mathbf{P}}(0) - 1$ of the energy dependence of the production amplitude and contribution to the diffraction slope B from the $\gamma^* \rightarrow V$ transition vertex, which is $\propto r_S^2$ and exhibits the $(Q^2 + m_V^2)$ scaling¹⁵, see fig. 2. This $(Q^2 + m_V^2)$ scaling formulated in 1974, has recently become a popular way of presenting the experimental data¹⁶. The theoretical calculations¹⁷ are based on the differential glue in the proton found in¹⁸

4 Shrinkage of the diffraction cone in hard diffraction

Gribov's shrinkage of the diffraction cone (DC) $B = B_0 + 2\alpha'_{\mathbf{P}} \log W^2$, quantized in terms of the slope $\alpha'_{\mathbf{P}}$ of the pomeron trajectory, is the salient feature of hadronic scattering, which derives from the Gribov-Feinberg-Chernavski diffusion in the impact parameter space. At this Symposium, we heard of the ZEUS finding¹⁶ of the shrinkage of the DC in $\gamma p \rightarrow J/\Psi p$ with the result $\alpha'_{eff} = 0.122 \pm 0.033(stat) + 0.018 - 0.032(syst) \text{ GeV}^{-2}$. Precisely such a shrinkage has been predicted in 1995 by Nikolaev, Zakharov and Zoller¹⁵ to persist within QCD even for hard processes.

First, in the usual approximation $\alpha_S = \text{const}$ the BFKL pomeron is the fixed cut in the complex- j plane¹⁹. Already in their first, 1975, publication on QCD pomeron, Kuraev, Lipatov and Fadin commented that incorporation of the asymptotic freedom splits the cut into a sequence of moving Regge poles¹⁹, see Lipatov²⁰ for more details. Within the CD approach, the Regge trajectories of these poles were calculated in^{15,22}. The CD cross section satisfies the CD BFKL equation²¹, $\partial\sigma(x, r)/\partial\log\frac{1}{x} = \mathcal{K} \otimes \sigma(x, r)$, which has the Regge solutions $\sigma_n(x, r) = \sigma_n(r)x^{-\Delta_n}$. The CD kernel $\mathcal{K}$ is related to the flux of Weizsäcker-Williams gluons around the $q\bar{q}$ dipole. The NZZ strategy was to evaluate α'_n from the energy dependence of $\lambda(x, r) = B\sigma(x, r)$, which satisfies the inhomogeneous equation $\partial\lambda(x, r)/\partial\log\frac{1}{x} - \mathcal{K} \otimes \lambda(x, r) = \mathcal{L} \otimes r^2\sigma(x, r)$, and has solutions $\lambda_n(x, r) = \sigma_n(x, r) \cdot \alpha'_n \log\frac{1}{x}$. Because $B = \frac{1}{2}\langle b^2 \rangle$ and the impact parameter b receives a contribution from the gluon- $q\bar{q}$ separation ρ , the inhomogeneous term $\mathcal{L} \otimes r^2\sigma(x, r)$ is driven by precisely the impact parameter diffusion of WW gluons. Because α'_n is driven by the inhomogeneous term, there is a manifest relationship between α'_n and the Gribov-Feinberg-Chernavski diffusion in the impact parameter space. Evidently, the dimensionfull quantity α'_n depends on the infrared regularization of QCD, within the specific regularization²¹ which has lead to an extremely successful description of the proton structure function (see²³ and references therein), for the rightmost hard BFKL pole we found $\alpha'_{\text{IP}} \approx 0.07 \text{ GeV}^{-2}$. The contribution from subleading BFKL poles was found to be still substantial at subasymptotic energy of HERA with the result $\alpha'_{eff} \approx 0.15 - 0.17 \text{ GeV}^{-2}$ ¹⁵, this prediction from 1995 agrees perfectly with the ZEUS finding.

5 Conclusions

The QCD theory of diffractive DIS is gradually coming of age. The fundamental $(Q^2 + m_V^2)$ scaling predicted in 1994 has finally been recognized by experimentalists. The shrinkage of the diffraction cone for hard photoproduction $\gamma p \rightarrow J/\Psi p$ discovered by the ZEUS collaboration is the single most important result. It shows that the BFKL pomeron is a (series of) moving pole(s) in the complex- j plane. The slope of the pomeron trajectory and the rate of shrinkage of the diffraction cone for hard photo- and electroproduction predicted in 1995 has been confirmed experimentally.

I'm grateful to the Symposium organizers, especially Profs. Liu Lian-shou and J. Crittenden, for the invitation to ISMD31. This work was partly supported by the INTAS grants 97-30494 and 00-00366.

References

1. N.N. Nikolaev, B.G. Zakharov and V.R. Zoller, *Z. Phys.* A351, 435 (1995)
2. N.N. Nikolaev and B.G. Zakharov, *Phys. Lett.* B55, 397 (1975)
3. E. Iancu, these proceedings.
4. J. Nemchik, N.N. Nikolaev and B.G. Zakharov, *Phys. Lett.* B341 (1994) 228; B.Z. Kopeliovich, J. Nemchik, N.N. Nikolaev and B.G. Zakharov, *Phys. Lett.* B309 (1993) 179; *Phys. Lett.* B324 (1994) 469
5. U. Maor, these proceedings; A. Gotsman, these proceedings.
6. N.N. Nikolaev and B.G. Zakharov, *Z. Phys.* C49, 607 (1991)
7. N.N. Nikolaev and B.G. Zakharov, *Phys. Lett.* B332, 184 (1996). .
8. N.N. Nikolaev and B.G. Zakharov, *Z. Phys.* C53, 331 (1992).
9. N.N. Nikolaev and B.G. Zakharov, *JETP* 78, 598 (1994); *Z. Phys.* C64 (1994) 631.
10. M. Genovese, N. Nikolaev and B. Zakharov, *JETP* 81 625 (1995); *JETP* 81, 633 (1995).
11. M. Genovese, N. Nikolaev and B. Zakharov, *Phys. Lett.* B378, 347 (1996); B380, 213 (1996).
12. C. Royon, L. Schoeffel, J. Bartels, H. Jung and R. Peschanski, *Phys. Rev. D* 63, 074004 (2001)
13. E. Elsen, these proceedings
14. H1 Collab: C. Adloff et al., *Eur. Phys. J.* C13 (2000) 371; ZEUS Collab: J. Breitweg et al.): *Eur. Phys. J.* C6, 603 (1999) and references therein
15. N.N. Nikolaev, B.G. Zakharov and V.R. Zoller, *Phys. Lett.* B366 (1996) 337; J. Nemchik, N.N. Nikolaev, E. Predazzi, B.G. Zakharov and V.R. Zoller, *J. Exp. Theor. Phys.* 86, 1054 (1998)
16. A. Savin, these proceedings
17. I.P. Ivanov & N.N. Nikolaev, 8th International Workshop on Deep Inelastic Scattering - DIS 2000, Liverpool, England, 25-30 April 2000, editors J.A. Gracey and T. Greenshaw, World Scientific, p. 636.
18. I.P. Ivanov and N.N. Nikolaev, *Phys. Atom. Nucl.* 64 (2001) 753; *Yad. Fiz.* 64 (2001) 813; *Phys. Rev. D*, in print.
19. V.S. Fadin, E.A. Kuraev and L.N. Lipatov *Phys. Lett.* B60 (1975) 50
20. L.N. Lipatov, *Sov. Phys. JETP* 63 (1986) 904.
21. N.N. Nikolaev, B.G. Zakharov and V.R. Zoller, *JETP Lett.* 59, 6 (1994)
22. N.N. Nikolaev, B.G. Zakharov and V.R. Zoller, *JETP Lett.* 66, 138 (1997), *Pisma Zh. Eksp. Teor. Fiz.* 66, 134 (1997)
23. N.N. Nikolaev, J. Speth and V.R. Zoller, *Phys. Lett.* B473, 157 (2000)